

**PATIENT INFORMATION LEAFLET**  
**GAVISCON STRAWBERRY FLAVOUR TABLETS**

**Sodium alginate**  
**Sodium hydrogen carbonate**  
**Calcium carbonate**

**Please read this leaflet carefully before you take this medicine.**  
**If you are not sure about anything ask your pharmacist or doctor.**

**What are Gaviscon Strawberry Flavour Tablets?**

Gaviscon Strawberry Flavour Tablets belong to a group of medicines called 'reflux suppressants', which form a protective layer on the top of the stomach contents to prevent stomach acid from escaping from the stomach where it works into the food pipe causing pain and discomfort.

**What are Gaviscon Strawberry Flavour Tablets used for?**

Gaviscon Strawberry Flavour Tablets are used for the treatment of symptoms of gastro-oesophageal reflux such as acid regurgitation, heartburn and indigestion, which can occur for example following meals or during pregnancy.

**Before taking Gaviscon Strawberry Flavour Tablets:**

**Do not take this product if:**

- you know you are allergic to any of the ingredients as very rarely difficulty in breathing and skin rashes have occurred (see Further information for a full list of ingredients).

**Take special care before treatment with Gaviscon Strawberry Flavour Tablets**

This medicine contains sodium (10.6mmol per four tablet dose) and calcium (3.2mmol per four tablet dose).

- If you have been advised to follow a diet restricted in either of these please consult your doctor.
- Please also talk to your doctor regarding these salt contents if you suffer or have suffered from significant kidney or heart disease, as certain salts could interfere with these diseases.

If you have phenylketonuria, note that the product is sweetened with aspartame, a source of phenylalanine.

Please consult your doctor if you know you have reduced amounts of gastric acid in your stomach, as this product may be less effective.

**Pregnancy and Breastfeeding**

You can take this product if you are pregnant or breastfeeding. As with all medicines, the treatment duration should be limited as much as possible.

0311223



### **Taking Other Medicines**

Do **not** take this product within two hours of taking other medicines by mouth as it can interfere with the action of some other medicines. This is especially important if you are taking antihistamines, antibiotics (tetracyclines and quinolones such as norfloxacin), iron preparations, antifungals such as ketoconazole, digoxin and beta-blockers (for heart conditions), neuroleptics (for mental illness), chloroquine (for malaria) and bisphosphonates (for osteoporosis).

### **How to take Gaviscon Strawberry Flavour Tablets:**

For oral administration. Chew thoroughly before swallowing.

**Adults, including the elderly and children 12 years and over:** When symptoms occur take two to four tablets after meals and at bedtime (up to four times a day).

**Children under 12 years:** Should only be taken on medical advice.

If you forget a dose it is not necessary to double the dose next time, just carry on taking as before.

If you take too much of this product you may feel bloated. It is unlikely to cause you any harm, but please consult your doctor or pharmacist.

### **After taking Gaviscon Strawberry Flavour Tablets:**

If symptoms persist after 7 days consult your doctor.

### **Possible side effects:**

Very rare (less than 1 in 10,000) chance of an allergic reaction to the ingredients. Symptoms of this may include skin rash, itching, difficulty breathing, dizziness, or swelling of the face, lips, tongue or throat. If you experience these or any other side-effects stop taking the product and consult your doctor immediately.

### **Storing your medicine:**

Do not use this product after the expiry date (EXP: month/year) shown.

**KEEP OUT OF THE REACH AND SIGHT OF CHILDREN.**

Do not store above 25°C. Store in the original package.

### **Further information:**

Gaviscon Strawberry Flavour Tablets are pale pink, circular, flat tablets with bevelled edges with the odour and flavour of strawberry.

Each chewable tablet contains 250mg sodium alginate, 133.5mg sodium hydrogen carbonate and 80mg calcium carbonate as the active ingredients. The other ingredients are Mannitol (E421), macrogol 20,000, aspartame (E951), magnesium stearate, xylitol and carmellose sodium, red iron oxide (E172), strawberry cream flavour.

Gaviscon Strawberry Flavour Tablets are available in pack sizes of 8, 16, 24, 32 tablets.

Manufacturer and PL Holder in the UK:

Reckitt Benckiser Healthcare (UK) Ltd., Hull, HU8 7DS.

PL 00063/0155.

Text Revised: June 2010.

0311223